

Durham Research Online

Deposited in DRO:

11 December 2018

Version of attached file:

Accepted Version

Peer-review status of attached file:

Peer-reviewed

Citation for published item:

Dalton, A.S. and Finkelstein, S.A. and Forman, S.L. and Barnett, P.J. and Pico, T. and Mitrovica, J.X. (2019) 'Was the Laurentide Ice Sheet significantly reduced during Marine Isotope Stage 3.', *Geology*, 47 (2). pp. 111-114.

Further information on publisher's website:

<https://doi.org/10.1130/G45335.1>

Publisher's copyright statement:

Additional information:

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

1 Was the Laurentide Ice Sheet significantly reduced during
2 Marine Isotope Stage 3?

3 April S. Dalton^{1,2}, Sarah A. Finkelstein¹, Steven L. Forman³, Peter J. Barnett⁴,
4 Tamara Pico⁵, and Jerry X. Mitrovica⁵

5 ¹*Department of Earth Sciences, University of Toronto, Toronto M5S 3B1, Canada*

6 ²*Department of Geography, Durham University, Durham DH1 3LE, United Kingdom*

7 ³*Department of Geosciences, Baylor University, Waco, Texas 76798, USA*

8 ⁴*Department of Earth Sciences, Laurentian University, Sudbury P3E 2C6, Canada*

9 ⁵*Department of Earth and Planetary Sciences, Harvard University, Cambridge,*
10 *Massachusetts 02138, USA*

11 **ABSTRACT**

12 Accurately reconstructing the paleogeography of the Laurentide Ice Sheet (LIS)
13 during Marine Isotope Stage 3 (MIS 3; ca. 57,000 to ca. 29,000 yr B.P.) is critical for
14 understanding glacial growth toward the Last Glacial Maximum (LGM), refining sea-
15 level histories and studying the Earth system response to rapid climate change events.
16 Here, we present a geochronological data set useful for testing hypotheses of global sea
17 level and refining ice sheet configuration through this interval. Data (n = 735) span the
18 entire MIS 3 interval and consist of ¹⁴C determinations (n = 651), cosmogenic exposure
19 ages (n = 52), and optically stimulated luminescence dates (n = 32). On that basis, we
20 hypothesize that the central region of the LIS underwent a dramatic reduction in ice from
21 ~52–40 ka. Key to this hypothesis are geological records at sites in the Hudson Bay

Lowlands that suggest a marine incursion and development of terrestrial landscapes. We show that these landscapes are consistent with recently published glacial isostatic adjustment predictions that include widespread deglaciation of the eastern (Labrador) sector of the LIS with ice build-up over the western (Keewatin) sector at 42 ka. Ice growth from this minimum toward the LGM is likely to have been rapid. The agreement between this data set and modeling predictions prompts the reassessment of key Late Pleistocene records, including Heinrich Events, loess deposition in the continental United States and sedimentological records from the Gulf of Mexico.

INTRODUCTION

The Laurentide Ice Sheet (LIS) was the predominant ice mass over North America through the Late Pleistocene (from ~125 to ~7 ka before present). Despite the importance of this ice sheet, its evolution is very poorly constrained prior to the Last Glacial Maximum (LGM; ~26 ka; Clark et al., 2009), especially during the interstadial of Marine Isotope Stage 3 (MIS 3; 59–27 ka; Lisiecki and Raymo, 2005). Refining ice sheet history through this interval will provide much-needed constraint on glacial growth toward the LGM; will aid in refining highly variable estimates of global mean sea level (GMSL; estimates range from –80m to –30m; Siddall et al., 2008) and; will offer a critical long-term perspective on the response of the Earth System to rapid climate events (e.g. Dansgaard-Oeschger and Heinrich events).

The most recent evaluation of geological data (via a synthesis of ~200 radiocarbon dates; Dyke et al., 2002) inferred that the LIS reached its MIS 3 minimum extent between 30–27 ka and was moderate in size, covering most of eastern and central Canada (Fig. 1). This ice configuration is often taken to represent the minimum extent

during the whole of MIS 3. However, since the work of Dyke et al. (2002), significantly more sites have been assigned to MIS 3. Notably, recent chronostratigraphic work in the Hudson Bay Lowlands, Canada (Dalton et al., 2016), lying at the geographic center of the LIS, suggests a significantly earlier and more pronounced interstadial minimum than hitherto recognized. Here, we (1) present an updated synthesis and evaluation of geological data for delineating the ice margin through the MIS 3 interval; (2) discuss the strength of geochronological data that support dramatic ice recession during MIS 3, and; (3) test the feasibility of this hypothesis by comparing geochronological data from the Hudson Bay Lowlands (a coastal plain where marine/terrestrial transitions are directly controlled by the interplay between ice loading, isostatic adjustment and sea level) with outputs from a recently published glacial isostatic adjustment (GIA) model for this time interval (ICE-PC2; Pico et al., 2017).

SYNTHESIS OF MIS 3 GEOLOGICAL DATA

The spatial and temporal extent of MIS 3 geochronological data¹ in the glaciated region ($n = 735$) has improved substantially since the last collective examination by Dyke et al. (2002). Available data now span all of MIS 3, with the majority of ages (56%) falling between 37.5–47.5 ka (Fig. 1). These sites document diverse and widespread ecosystems during MIS 3 (e.g., boreal forest, peatlands) in regions that were later overrun by ice during the LGM. Samples consist of radiocarbon determinations (88.6%, $n = 651$), cosmogenic exposure ages (7.1%; $n = 52$) and luminescence dates (4.4%; $n = 32$). Preservation of MIS 3 sediments is largely in geological contexts that offered protection from LGM glacial advance, such as river valleys, coastal cliffs and deep lacustrine environments. For this reason, pre-LGM stratigraphic records are rarely preserved on the

Canadian Shield, a granitic geological unit covering a large swath of Canada (Fig. 1; gray shading). Geochronological data from much of the glaciated region are now of sufficient quality to track landscape evolution through MIS 3 at millennial-scales (e.g. Hughes et al., 2016), critical for testing the validity of glaciological models that have been developed for this interval (e.g., Stokes et al., 2012).

EVIDENCE SUPPORTING DRAMATIC REDUCTION OF THE LAURENTIDE ICE SHEET

Geochronological data from the Hudson Bay area (n=35) suggest that the LIS underwent large-scale deglaciation during MIS 3. Data from this region (^{14}C and optically stimulated luminescence dating of marine and fluvial strata) support a marine incursion between 52 and 42 ka, along with the subsequent development of terrestrial landscapes during the interval of 48–40 ka (Dalton et al., 2016). Deglaciation in this central region was first hypothesized in the 1980s (Andrews et al., 1983; Dredge and Thorleifson, 1987). However, at that time, this hypothesis was largely dismissed based on suspected inaccuracy of chronological techniques, especially the reliance on ^{14}C dating of shells and amino acid dating. Now, multiple radiocarbon age attempts per site (on wood, where possible), along with confirmatory OSL dating suggests that the MIS 3 age assignment may indeed be correct.

The hypothesis of a significantly reduced ice sheet can be tested by comparing geochronological data from key areas to predictions from a GIA model. In this regard, we note that available geochronological data from the Hudson Bay Lowlands offer a good fit with a glacial-isostatic adjustment simulation forced by ICE-PC2, an ice loading history with widespread deglaciation of the eastern sector of the LIS during MIS 3 (to

accommodate high sea level markers of this age along the east coast of the United States) with ice build-up preferentially over Keewatin at 42 ka (Fig. 2; Pico et al., 2017). Notably, the marine-estuarine deposits on the Severn River dated to 42–52 ka (Dalton et al., 2016; Forman et al., 1987), align well with the simulated paleo-shoreline in that region. Further, terrestrial sites dated to this interval (11-PJB-186, 11-PJB-020, 12-PJB-007; Dalton et al., 2016) are all located at sub-aerial elevations in the simulation (Fig. 2A). The discrepancy between the simulated location of the shoreline and the marine sites suggests that adjustments to the ice load may be necessary to accommodate all data, within error. The ICE-PC2 simulation is consistent with mid-MIS 3 dates on non-glacial deposits in the Hudson Bay Lowlands and adopts a mid-MIS 3 relative sea-level highstand at 44 ka with a GMSL value of –38 m (Fig. 2B; Pico et al., 2016; additional details in Data Repository¹).

DISCUSSION

In support of our hypothesis, insolation in the Northern Hemisphere during MIS 3 was at its most stable point for the entire Late Pleistocene, and slightly higher than present-day (Fig. 3A; Berger and Loutre, 1991) which may suggest that climate forcing was sufficient to drive a prolonged, dynamic reduction of continental ice. This insolation promoted a pronounced recession in the Fennoscandian Ice Sheet (Helmens and Engels, 2010). Fluctuations in global atmospheric methane during MIS 3 (Fig. 3B; Loulergue et al., 2008) could be partly explained by the development of northern peatlands in the Hudson Bay Lowlands during that time (Dalton et al., 2017). Also noteworthy are perturbations in the $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ record from Crevice Cave (Dorale et al., 1998) and Devils Hole (Landwehr et al., 2011) during MIS 3 (Figs. 3C–E) that suggest pronounced

changes in North American precipitation regimes and vegetation during that interval, possibly as a result of reduced and/or highly variable continental ice. Our hypothesis is also supported by geomorphological records (striations and glacial lineations; Kleman et al., 2010) suggesting the eastern and western sectors of the LIS were independent for an extended period prior to the LGM, which some numerical modeling has failed to reproduce (Stokes et al., 2012) largely due to a lack of geochronological constraints.

An important corollary of our hypothesis is that it prompts the reassessment of key Late Pleistocene records. Reduced continental ice likely caused meltwater to flow northward toward the Arctic Ocean during some of MIS 3. Thus, any meltwater-based control over Dansgaard-Oeschger Events through this interval (Fig. 3F) must have been dominated by northward drainage perturbations (as opposed to alternating with the Gulf of Mexico; Clark et al., 2001). Geological records from the Mississippi River that have been widely assigned to ice advance into the mid-continent may also require reinterpretation; these include the widespread Roxana Silt loess deposit (~60–30 ka interval; see Forman and Pierson, 2002, and references therein), pronounced fluvial aggradation along the Mississippi River (Rittenour et al., 2007) and sedimentological shifts in the Gulf of Mexico (Sionneau et al., 2013). Possible non-glacial explanations for the increased sediment flux and aggradation include changes in catchment vegetation, seasonality of rainfall and variable hydrology rather than a direct meltwater signature (Leigh et al., 2004). Our hypothesis also suggests that full glaciation of Hudson Bay is not necessarily a prerequisite for Heinrich Events given that both Heinrich Event 5 (H5; 50–47 ka) and 4 (H4; 40.2–38.3 ka) (Sanchez Goñi and Harrison, 2010) took place around the time interval of interest. It is possible that Baffin Island, Southampton Island

or Labrador could have been active contributors to ice rafting events in the north-eastern LIS (Hefter et al., 2017; Roy et al., 2009). In this case, adjustments to the modeled Labrador sector may be needed (Fig. 2).

Growth of the LIS from this receded position was likely very rapid. Using a combination of geological constraints and numerical modelling, recent studies have demonstrated that the Labrador sector may have expanded >1000 km southward by ~39–37 ka (Carlson et al., 2018; Pico et al., 2018a,b). We acknowledge that continued work is needed to understand the evolution of the MIS 3 landscape, study the impact of reduced continental ice on global systems, and delineate precise ice growth toward the LGM. Nevertheless, the agreement between available geological data, GMSL and geophysical modelling supports a significant reduction of the LIS during MIS 3.

ACKNOWLEDGMENTS

This research was funded by grants to SAF from the Natural Sciences and Engineering Research Council (Canada) and the Ontario Geological Survey; and to ASD from the Northern Scientific Training Program, the University of Toronto Centre for Global Change Science and the Ontario Graduate Scholarship. We acknowledge the support of the National Science Foundation to TP and support from Harvard University to TP and JXM. We thank Chris Stokes for insightful comments, Art Dyke for providing some chronology data, John Andrews for conversations about Heinrich Events, and three anonymous reviewers for constructive comments.

REFERENCES CITED

- 158 Andrews, J.T., Shilts, W.W., and Miller, G.H., 1983, Multiple deglaciations of the
159 Hudson Bay Lowlands, Canada, since deposition of the Missinaibi (Last-
160 Interglacial?): Formation: Quaternary Research, v. 19, p. 18–37.
- 161 Berger, A., and Loutre, M.F., 1991, Insolation values for the climate of the last 10 million
162 years: Quaternary Science Reviews, v. 10, p. 297–317, [https://doi.org/10.1016/0277-](https://doi.org/10.1016/0277-3791(91)90033-Q)
163 3791(91)90033-Q.
- 164 Carlson, A. E., Tarasov, L., and Pico, T., 2018, Rapid Laurentide ice-sheet advance
165 towards southern last glacial maximum limit during marine isotope stage 3:
166 Quaternary Science Reviews, v. 196, p. 118-123,
167 <https://doi.org/10.1016/j.quascirev.2018.07.039>
- 168 Clark, P. U., Dyke, A. S., Shakun, J. D., Carlson, A. E., Clark, J., Wohlfarth, B.,
169 Mitrovica, J. X., Hostetler, S. W., and McCabe, A. M., 2009, The Last Glacial
170 Maximum: Science, v. 325, p. 710–714, <https://doi.org/10.1126/science.1172873>.
- 171 Clark, P.U., Marshall, S.J., Clarke, G.K.C., Hostetler, S.W., Licciardi, J.M., and Teller,
172 J.T., 2001, Freshwater forcing of abrupt climate change during the last glaciation:
173 Science, v. 293, p. 283–287, <https://doi.org/10.1126/science.1062517>.
- 174 Dalton, A.S., Finkelstein, S.A., Barnett, P.J., and Forman, S.L., 2016, Constraining the
175 Late Pleistocene history of the Laurentide Ice Sheet by dating the Missinaibi
176 Formation, Hudson Bay Lowlands, Canada: Quaternary Science Reviews, v. 146,
177 p. 288–299, <https://doi.org/10.1016/j.quascirev.2016.06.015>.
- 178 Dalton, A.S., Väliranta, M., Barnett, P.J., and Finkelstein, S.A., 2017, Pollen and
179 macrofossil-inferred paleoclimate at the Ridge Site, Hudson Bay Lowlands, Canada:
180 Evidence for a dry climate and significant recession of the Laurentide Ice Sheet

- 181 during Marine Isotope Stage 3: Boreas, v. 46, p. 388–401,
182 <https://doi.org/10.1111/bor.12218>.
- 183 Dorale, J.A., Edwards, R.L., Ito, E., and González, L.A., 1998, Climate and Vegetation
184 History of the Midcontinent from 75 to 25 ka: A Speleothem Record from Crevice
185 Cave, Missouri, USA: *Science*, v. 282, p. 1871–1874,
186 <https://doi.org/10.1126/science.282.5395.1871>.
- 187 Dredge, L.A., and Thorleifson, L.H., 1987, The Middle Wisconsinan History of the
188 Laurentide Ice Sheet: *Géographie physique et Quaternaire*, v. 41, p. 215–235,
189 <https://doi.org/10.7202/032680ar>.
- 190 Dyke, A.S., 2004, An outline of North American deglaciation with emphasis on central
191 and northern Canada, *in* Ehlers, J., and Gibbard, P. L., eds., *Quaternary Glaciations -*
192 *Extent and Chronology, Part II: Amsterdam, the Netherlands, Elsevier*, p. 373–424,
193 [https://doi.org/10.1016/S1571-0866\(04\)80209-4](https://doi.org/10.1016/S1571-0866(04)80209-4).
- 194 Dyke, A.S., Andrews, J.T., Clark, P.U., England, J.H., Miller, G.H., Shaw, J., and
195 Veillette, J.J., 2002, The Laurentide and Innuitian ice sheets during the Last Glacial
196 Maximum: *Quaternary Science Reviews*, v. 21, p. 9–31,
197 [https://doi.org/10.1016/S0277-3791\(01\)00095-6](https://doi.org/10.1016/S0277-3791(01)00095-6).
- 198 Forman, S.L., and Pierson, J., 2002, Late Pleistocene luminescence chronology of loess
199 deposition in the Missouri and Mississippi river valleys, United States:
200 *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 186, p. 25–46,
201 [https://doi.org/10.1016/S0031-0182\(02\)00440-6](https://doi.org/10.1016/S0031-0182(02)00440-6).
- 202 Forman, S.L., Wintle, A.G., Thorleifson, L.H., and Wyatt, P.H., 1987,
203 Thermoluminescence properties and age estimates for Quaternary raised marine

- 204 sediments, Hudson Bay Lowland, Canada: Canadian Journal of Earth Sciences,
205 v. 24, p. 2405–2411, <https://doi.org/10.1139/e87-226>.
- 206 Hefter, J., Naafs, B.D.A., and Zhang, S., 2017, Tracing the source of ancient reworked
207 organic matter delivered to the North Atlantic Ocean during Heinrich Events:
208 *Geochimica et Cosmochimica Acta*, v. 205, p. 211–225,
209 <https://doi.org/10.1016/j.gca.2017.02.008>.
- 210 Helmens, K., and Engels, S., 2010, Ice-free conditions in eastern Fennoscandia during
211 early Marine Isotope Stage 3: lacustrine records: *Boreas*, v. 39, p. 399–409,
212 <https://doi.org/10.1111/j.1502-3885.2010.00142.x>.
- 213 Hughes, A. L. C., Gyllencreutz, R., Lohne, Ø. S., Mangerud, J., and Svendsen, J. I., 2016,
214 The last Eurasian ice sheets – a chronological database and time-slice reconstruction,
215 DATED-1: *Boreas*, v. 45, p. 1–45, <https://doi.org/10.1111/bor.12142>.
- 216 Kleman, J., Jansson, K., De Angelis, H., Stroeve, A.P., Hättestrand, C., Alm, G., and
217 Glasser, N., 2010, North American Ice Sheet build-up during the last glacial cycle,
218 115–21kyr: *Quaternary Science Reviews*, v. 29, p. 2036–2051,
219 <https://doi.org/10.1016/j.quascirev.2010.04.021>.
- 220 Landwehr, J.M., Sharp, W.D., Coplen, T.B., Ludwig, K.R., and Winograd, I.J., 2011, The
221 chronology for the $\delta^{18}\text{O}$ record from Devils Hole, Nevada, extended into the mid-
222 Holocene: U.S. Geological Survey Open-File Report 2011–1082, 5 p.
- 223 Leigh, D., Srivastava, P., and Brook, G., 2004, Late Pleistocene braided rivers of the
224 Atlantic Coastal Plain, USA: *Quaternary Science Reviews*, v. 23, p. 65–84,
225 [https://doi.org/10.1016/S0277-3791\(03\)00221-X](https://doi.org/10.1016/S0277-3791(03)00221-X).

- 226 Lisiecki, L.E., and Raymo, M.E., 2005, A Pliocene-Pleistocene stack of 57 globally
227 distributed benthic $\delta^{18}\text{O}$ records: *Paleoceanography*, v. 20, p. PA1003.
- 228 Loulergue, L., Schilt, A., Spahni, R., Masson-Delmotte, V., Blunier, T., Lemieux, B.,
229 Barnola, J.-M., Raynaud, D., Stocker, T.F., and Chappellaz, J., 2008, Orbital and
230 millennial-scale features of atmospheric CH_4 over the past 800,000 years: *Nature*,
231 v. 453, p. 383–386, <https://doi.org/10.1038/nature06950>.
- 232 North Greenland Ice Core Project members, 2004, High-resolution record of Northern
233 Hemisphere climate extending into the last interglacial period: *Nature*, v. 431,
234 p. 147–151, <https://doi.org/10.1038/nature02805>.
- 235 Pico, T., Birch, L., Weisenberg, J., and Mitrovica, J. X., 2018b, Refining the Laurentide
236 Ice Sheet at Marine Isotope Stage 3: A data-based approach combining glacial
237 isostatic simulations with a dynamic ice model: *Quaternary Science Reviews*, v. 195,
238 p. 171–179, <https://doi.org/10.1016/j.quascirev.2018.07.023>.
- 239 Pico, T., Creveling, J. R., and Mitrovica, J. X., 2017, Sea-Level Records from the U.S.
240 Mid-Atlantic Constrain Laurentide Ice Sheet Extent During Marine Isotope Stage 3:
241 *Nature Communications*, v. 8, p. 15612 / DOI:10.1038/ncomms15612.
- 242 Pico, T., Mitrovica, J.X., Braun, J., and Ferrier, K.L., 2018a, Glacial isostatic adjustment
243 deflects the path of the ancestral Hudson River: *Geology*, v. 46, p. 591–594,
244 <https://doi.org/10.1130/G40221.1>.
- 245 Pico, T., Mitrovica, J.X., Ferrier, K.L., and Braun, J., 2016, Global ice volume during
246 MIS 3 inferred from a sea-level analysis of sedimentary core records in the Yellow
247 River Delta: *Quaternary Science Reviews*, v. 152, p. 72–79,
248 <https://doi.org/10.1016/j.quascirev.2016.09.012>.

- 249 Rittenour, T.M., Blum, M.D., and Goble, R.J., 2007, Fluvial evolution of the lower
250 Mississippi River valley during the last 100 k.y. glacial cycle: Response to glaciation
251 and sea-level change: Geological Society of America Bulletin, v. 119, p. 586–608,
252 <https://doi.org/10.1130/B25934.1>.
- 253 Roy, M., Hemming, S.R., and Parent, M., 2009, Sediment sources of northern Québec
254 and Labrador glacial deposits and the northeastern sector of the Laurentide Ice Sheet
255 during ice-rafting events of the last glacial cycle: Quaternary Science Reviews, v. 28,
256 p. 3236–3245, <https://doi.org/10.1016/j.quascirev.2009.08.008>.
- 257 Sanchez Goñi, M.F., and Harrison, S.P., 2010, Millennial-scale climate variability and
258 vegetation changes during the Last Glacial: Concepts and terminology: Quaternary
259 Science Reviews, v. 29, p. 2823–2827,
260 <https://doi.org/10.1016/j.quascirev.2009.11.014>.
- 261 Siddall, M., Rohling, E.J., Thompson, W.G., and Waelbroeck, C., 2008, Marine Isotope
262 Stage 3 sea level fluctuations: Data synthesis and new outlook: Reviews of
263 Geophysics, v. 46, p. RG4003, <https://doi.org/10.1029/2007RG000226>.
- 264 Sionneau, T., Bout-Roumazeilles, V., Meunier, G., Kissel, C., Flower, B.P., Bory, A.,
265 and Tribouvillard, N., 2013, Atmospheric re-organization during Marine Isotope Stage
266 3 over the North American continent: Sedimentological and mineralogical evidence
267 from the Gulf of Mexico: Quaternary Science Reviews, v. 81, p. 62–73,
268 <https://doi.org/10.1016/j.quascirev.2013.10.002>.
- 269 Stokes, C.R., Tarasov, L., and Dyke, A.S., 2012, Dynamics of the North American Ice
270 Sheet Complex during its inception and build-up to the Last Glacial Maximum:

Quaternary Science Reviews, v. 50, p. 86–104,

<https://doi.org/10.1016/j.quascirev.2012.07.009>.

FIGURE CAPTIONS

Figure 1. Map showing the glacial outline of the Laurentide Ice Sheet at 30–27 ka (after Dyke et al., 2002), overlaid with currently available geological data from Marine Isotope Stage 3 (MIS 3; $n = 735$). Note the large number of dates that lie inside the MIS 3 extent (e.g., Hudson Bay Lowlands), which likely indicate an earlier and significantly more pronounced ice reduction prior to the 30–27 ka interval. Shaded region is the Canadian Shield. Last Glacial Maximum (LGM) ice extent after Dyke (2004). Some sites have multiple ages that overlap on this plot; all geological data are available in Table DR1. Inset figure shows the age distribution of chronology data spanning MIS 3 with data binned into 2500-year increments. NB: data plotted here are not necessarily in conflict with the work of Dyke et al., (2002) since the ice perimeter in the 2002 study was intended to represent only the 30 –27 ka interval.

Figure 2. Predicted North American topography at 42 ka using ice history ICE-PC2, overlaid with available geological data from the Hudson Bay Lowlands (colored as in Fig. 1). Note the agreement between available geochronological data and the numerical simulation, which supports the hypothesis of reduced ice cover during this interval. Present-day coastline shown by black contours. Shoreline at 42 ka (0 m contour) shown by gray outline. A: Zoom into local topography of the Hudson Bay Lowlands. B: Global mean sea level change adopted in ice history ICE-PC2 (Pico et al., 2017).

294 Figure 3. Paleoclimate and orbital parameters spanning the Late Pleistocene (100 ka to
295 present-day). A: July insolation at 60°N after Berger and Loutre (1991). B: Atmospheric
296 methane estimates from the EPICA Dome C ice core (Loulergue et al., 2008). C–D:
297 Carbon and oxygen isotope data from Crevice Cave (Dorale et al., 1998). E–F: Oxygen
298 isotope data from Devils Hole speleothem (Landwehr et al., 2011) and a northern
299 Greenland ice core (North Greenland Ice Core Project members, 2004). Arrows indicate
300 Heinrich Events (Sanchez Goñi and Harrison, 2010).

301

302 1GSA Data Repository item 2018xxx, table of geochronological data (n=735) and
303 description of the geophysical model, is available online at
304 <http://www.geosociety.org/datarepository/2018/>, or on request from
305 editing@geosociety.org.




Figure 1

Dalton et al. (in press)




Figure 2

Dalton et al. (in press)

Figure 3
Dalton et al. (in press)

